

2003 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS (Form B)

CHEMISTRY

Section II

(Total time—90 minutes)

Part A

Time—40 minutes

YOU MAY USE YOUR CALCULATOR FOR PART A.


CLEARLY SHOW THE METHOD USED AND THE STEPS INVOLVED IN ARRIVING AT YOUR ANSWERS. It is to your advantage to do this, since you may obtain partial credit if you do and you will receive little or no credit if you do not. Attention should be paid to significant figures.

Be sure to write all your answers to the questions on the lined pages following each question in the booklet with the goldenrod cover. Do NOT write your answers on the lavender insert.

Answer Question 1 below. The Section II score weighting for this question is 20 percent.



1. After a 1.0 mole sample of $\text{HI}(g)$ is placed into an evacuated 1.0 L container at 700. K, the reaction represented above occurs. The concentration of $\text{HI}(g)$ as a function of time is shown below.



- (a) Write the expression for the equilibrium constant, K_c , for the reaction.
- (b) What is $[\text{HI}]$ at equilibrium?

2003 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS (Form B)

- (c) Determine the equilibrium concentrations of $\text{H}_2(g)$ and $\text{I}_2(g)$.
- (d) On the graph above, make a sketch that shows how the concentration of $\text{H}_2(g)$ changes as a function of time.
- (e) Calculate the value of the following equilibrium constants at 700. K.
- (i) K_c
 - (ii) K_p
- (f) At 1,000 K, the value of K_c for the reaction is 2.6×10^{-2} . In an experiment, 0.75 mole of $\text{HI}(g)$, 0.10 mole of $\text{H}_2(g)$, and 0.50 mole of $\text{I}_2(g)$ are placed in a 1.0 L container and allowed to reach equilibrium at 1,000 K. Determine whether the equilibrium concentration of $\text{HI}(g)$ will be greater than, equal to, or less than the initial concentration of $\text{HI}(g)$. Justify your answer.